



## King's Research Portal

DOI:

[10.1016/j.jpsychires.2018.06.019](https://doi.org/10.1016/j.jpsychires.2018.06.019)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Taylor, C. L., Broadbent, M., Khondoker, M., Stewart, R., & Howard, L. M. (2018). Predictors of severe relapse in pregnant women with psychotic or bipolar disorders. *Journal of psychiatric research*, 104, 100-107.  
<https://doi.org/10.1016/j.jpsychires.2018.06.019>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Accepted Manuscript

Predictors of severe relapse in pregnant women with psychotic or bipolar disorders

Clare L. Taylor, Matthew Broadbent, Mizanur Khondoker, Robert Stewart, Louise M. Howard



PII: S0022-3956(17)31407-3

DOI: [10.1016/j.jpsychires.2018.06.019](https://doi.org/10.1016/j.jpsychires.2018.06.019)

Reference: PIAT 3410

To appear in: *Journal of Psychiatric Research*

Received Date: 8 February 2018

Revised Date: 18 June 2018

Accepted Date: 29 June 2018

Please cite this article as: Taylor CL, Broadbent M, Khondoker M, Stewart R, Howard LM, Predictors of severe relapse in pregnant women with psychotic or bipolar disorders, *Journal of Psychiatric Research* (2018), doi: 10.1016/j.jpsychires.2018.06.019.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title: Predictors of severe relapse in pregnant women with psychotic or bipolar disorders**

Clare L Taylor<sup>a\*</sup>, Matthew Broadbent<sup>b</sup>, Mizanur Khondoker<sup>c</sup>, Robert Stewart<sup>db</sup>, Louise M Howard<sup>ab</sup>

<sup>a</sup>Section of Women's Mental Health, Health Service and Population Research Department, Institute of Psychiatry, King's College London, UK

<sup>b</sup>South London and Maudsley NHS Foundation Trust, London, UK

<sup>c</sup>University of East Anglia, Norwich Medical School, Norwich Research Park, Norwich, UK

<sup>d</sup>Psychological Medicine Department, Institute of Psychiatry, King's College London, UK

**\*Corresponding author:** Clare L Taylor, Section of Women's Mental Health, Department of Health Services and Population Research, PO31 Institute of Psychiatry, Psychology & Neuroscience, King's College London, De Crespigny Park, London, SE5 8AF, UK

Author's email addresses: clare.l.taylor@kcl.ac.uk, matthew.broadbent@kcl.ac.uk, m.khondoker@uea.ac.uk, Robert.stewart@kcl.ac.uk

**Number of tables and figures:** - 1 figure, 4 tables, 7 supplementary tables

**Word count excluding abstract, tables, references and supplementary material:** - 4,384

**Abstract**

Pregnancy in women with severe mental illness is associated with adverse outcomes for mother and infant. There are limited data on prevalence and predictors of relapse in pregnancy. A historical cohort study using anonymised comprehensive electronic health records from secondary mental health care linked with national maternity data was carried out. Women with a history of serious mental illness who were pregnant (2007-2011), and in remission at the start of pregnancy, were studied; severe relapse was defined as admission to acute care or self-harm. Predictors of relapse were analysed using random effects logistic regression to account for repeated measures in women with more than one pregnancy in the study period. In 454 pregnancies (389 women) there were 58 (24%) relapses in women with non-affective psychoses and 25 (12%) in women with affective psychotic or bipolar disorders. Independent predictors of relapse included non-affective psychosis (adjusted OR=2.03; 95% CI=1.16-3.54), number of recent admissions (1.37; 1.03-1.84), recent self-harm (2.24; 1.15-4.34), substance use (2.15; 1.13-4.08), smoking (2.52; 1.26-5.02) and non-white ethnicity (black ethnicity: 2.37; 1.23,4.57, mixed/other ethnicity: 2.94; 1.32,6.56). Women on no regular medication throughout first trimester were also at greater risk of relapse in pregnancy (1.99; 1.05-3.75). There was no interaction between severity of illness and medication status as relapse predictors. Therefore, women with non-affective psychosis and higher number of recent acute admissions are at significant risk of severe relapse in pregnancy. Continuation of medication in women with severe mental illness who become pregnant may be protective.

**Keywords:** psychosis; bipolar; pregnancy; relapse; medication

## **Introduction**

Relapse of serious mental illness (SMI; schizophrenia, related delusional disorders, affective psychoses and bipolar disorder) in the perinatal period is potentially devastating for a woman and her unborn baby. It may impact on her ability to care for herself, and rarely but tragically result in custody loss of children, suicide or infanticide (Jones et al, 2014). International confidential enquiries into maternal deaths highlight psychiatric illness as a leading cause (Austin et al, 2007; Cantwell et al, 2011; Khalifeh et al, 2016).

Studies have consistently reported increased risk of relapse of psychosis in the early postpartum, with risk factors including history of bipolar disorder or post-partum psychosis (Jones et al, 2014; Munk-Olsen et al, 2009; Wesseloo et al, 2015). There are fewer data on the course of SMI during pregnancy, particularly for schizophrenia. Recently reported data indicated a 12% rate of hospitalisation for women with schizophrenia in pregnancy (Rochon-Terry et al, 2016). For bipolar disorder some studies estimate around 8-18% relapse in pregnancy (Di Florio et al, 2013; Grof et al, 2000), but tertiary perinatal mental health clinics have reported substantially higher rates (71%), particularly in women who discontinue medication (86%) and those with more severe illness (Viguera et al, 2007). Varying definitions of relapse may explain inconsistencies; some use hospital admission and others use clinical or research interviews to determine symptomatology, which identify less severe relapses.

We aimed to investigate risk and predictors of severe relapse in pregnancy in a cohort of women with SMI. We hypothesised that: 1) diagnosis and markers of severity (non-affective SMI, recent admissions) and 2) first trimester medication changes (switching or discontinuation) would be

independently associated with relapse. We also explored interactions between diagnosis, admissions in two years before pregnancy and domestic violence (DV) with medication.

## **Methods**

### **Data Source**

Data came from the South London and Maudsley (SLaM) NIHR Biomedical Research Centre Clinical Record Interactive Search (CRIS) (Stewart et al, 2009), a large resource of comprehensive secondary mental health records from over 300,000 patients. SLaM serves a catchment of around 1.2 million residents from four London boroughs, and provides some tertiary services. CRIS allows searching and retrieval of anonymised clinical records within robust technical and procedural safeguards (Fernandes et al, 2013). Data can be extracted from free text (e.g. case notes and correspondence), using targeted keyword searches. Structured data include socio-demographics (including ethnicity), year and month of birth, and referrals data including admissions to SLAM and movements within SLaM services such as wards and community teams. Several natural language processing (NLP) applications on a General Architecture for Text Engineering (GATE) platform (Cunningham et al, 2013) have been developed which derive structured data from free text and are validated against manual annotations. These include diagnosis and medication (Perera et al, 2016). CRIS was approved for research by Oxfordshire Research Ethics Committee C (08/H0606/71+5), and is linked to Hospital Episode Statistics: national data for inpatient, outpatient and accident and emergency attendances in England including maternity data.

### **Study population**

A historic cohort was assembled of women pregnant from 2007-2011 whose pregnancies continued beyond first trimester. We used maternity Hospital Episode Statistics (HES) indicating delivery of a baby linked to CRIS data indicating women with SMI, using ICD-10 diagnoses F20, F22, F23, F25, F28, F29 (schizophrenia and related disorders, schizoaffective disorders and delusional disorders), F30, F31 (mania and bipolar disorders), F32.3, F33.3 (psychotic depression), F53.1 (puerperal psychosis). CRIS data is complete from 2006 and at the time of data extraction, HES data was available up to the end of 2011. HES data indicated the end date of pregnancy. In order to collect time dependent

variables, trimesters of pregnancy were dated using a modified validated algorithm (Li et al, 2013; Taylor et al, 2015). CRIS data captured socio-demographic and clinical details regarding women's mental health. In order to be able to capture clinical details regarding women's mental health in the perinatal period, we included women under SLAM care (therefore on CRIS) at any point from 6 months before to 6 weeks after the HES delivery episode with SMI diagnoses during the preconception period so. We excluded women whose illness was not in remission at the start of pregnancy, determined by admission to acute mental health care (inpatient or home treatment) in the three months before the start date of pregnancy.

### Measures

#### *Severe relapse*

The first severe relapse of SMI (hereafter termed relapse) occurring during pregnancy was measured. This has previously been defined as hospitalisation (Wesseloo et al, 2015). We defined it as admission to mental health inpatient care, referral for home treatment (services for acute mental health crisis in the UK, involving frequent home visits (Johnson et al, 2008)) or an episode of self-harm (see (Taylor et al, 2016)) so that all significant relapses were included. Admissions/ referrals were extracted from structured CRIS data or HES inpatient admissions to mental health trusts outside SLAM. Self-harm during pregnancy was extracted using free text search terms previously validated in CRIS (Polling et al, 2015; Taylor et al, 2016).

#### *Socio-demographic characteristics for index (first) pregnancy*

Recorded ethnicity was categorised into 'Black African/Caribbean/other', 'White British/other', and 'Mixed, Asian or other/ not stated'. Neighbourhood-level deprivation score was extracted closest in date to the beginning of pregnancy. These give a summary measure of socioeconomic status at the level of the lower super output area (LSOA): a geographic unit containing a minimum of 1000 residents and 400 households with an average nationally of around 1500 residents (Noble



et al, 2007). Manual searches of CRIS free text using piloted terms established number of children prior to pregnancy, relationship status in pregnancy, history of childhood abuse and DV before/during pregnancy and family history of mental illness. Recorded smoking in pregnancy was extracted by manual free text searches using terms (Wu et al, 2013).

#### *Mental disorder characteristics and severity at the beginning of pregnancy*

Diagnosis recorded at the start of pregnancy, extracted from CRIS structured fields and free text using an NLP application validated for dementia (Perera et al, 2016), was categorised into non-affective and affective. Non-affective SMI comprised schizophrenia, related psychoses, delusional disorders and schizoaffective disorders. Affective SMI comprised bipolar disorder, psychotic depression and history of postpartum psychosis. Where diagnosis was unclear, correspondence was scrutinised by a researcher and clinician.

Number of days in acute mental health care in the two years before pregnancy was extracted from CRIS (supplemented with HES for outside-SLaM care (Taylor et al, 2015)) and spells of acute care defined as periods of treatment in acute care (inpatient or home treatment) with at most 7 days between discharge and readmission. Time since last major episode was calculated from the beginning of the most recent 'spell' of acute care in the 2 years prior to conception. Detention under the Mental Health Act (involuntary admission) in the 2 years before pregnancy was extracted from structured fields and free text.

Manual free text searches of CRIS ascertained recorded alcohol and drug use in pregnancy within the previous 2 years, supplemented by substance misuse diagnoses from CRIS structured fields and an NLP application. Baseline level of functioning was estimated from the highest total adjusted score recorded in the two years before pregnancy from the Health of the Nation Outcome Scale (HoNOS), a routinely collected 12-item measure in UK mental health services of health and social functioning of people with severe mental illness (Wing et al, 1998) stored in CRIS.

*Medication exposure in the first trimester of pregnancy*

Medication exposures were extracted from free text in CRIS, primarily using an NLP application (validated for a number of antipsychotic agents (Kadra et al, 2015)) to extract structured indicators describing medication during the first trimester, supplemented by manual text searches. The clinical documents were recalled and manually coded for information on medications not already validated. Stops, starts and switches in regular antipsychotics, mood stabilisers and antidepressants were noted, including whether stops occurred abruptly (over 1-14 days) or gradually (>14 days). Where speed of discontinuation was not specified, women were coded as stopping abruptly. Recorded non-adherence to medication was categorised if there was written concern about this. Two researchers rated medication, with consensus meetings to resolve uncertainties; a consecutive (26 pregnancies) were rated for identification of antipsychotic, mood stabiliser and antidepressant with inter-rater agreements of 97%. For the main analysis, medication was categorised as 'stopped or switched' if a woman had a stop and/or switch in a regular medication, 'continued' if a woman continued a regular medication all the way through the first trimester and did not stop or switch a regular medication, and 'no medication at start of pregnancy' if a woman was not on any regular medication at the start of 1<sup>st</sup> trimester. Further detail including exposures to individual medications and medication types in 3 months before pregnancy and 1<sup>st</sup> trimester is provided elsewhere (Taylor et al, 2015).

*Data analysis*

Data were analysed using STATA 12. After initial descriptive analyses of women, subsequent analyses were carried out by pregnancy. Incidence and timing (by trimester) of relapses during pregnancy were reported for the whole sample and stratified by baseline diagnosis. To account for non-independent data in women who contributed more than one pregnancy during the study period, all subsequent statistical analyses were carried out using random effects logistic regression. Only one indicator of acute mental health care before pregnancy – number of admissions – was used

in multivariable analyses following  $\chi^2$  tests for association between them to assess collinearity. We used purposeful selection of variables (Bursac et al, 2008), and predictors with  $p \leq 0.2$  from unadjusted analyses were added simultaneously to the model and unadjusted and fully adjusted effect sizes compared. Predictors with a fully adjusted p-value of  $>0.5$  were removed from the final model. Given our main hypothesis around medication; we ran post-hoc analyses using  $\chi^2$  tests to assess association of covariates with the medication variable using the 3 category medication variable as an outcome and random effect logistic regression to look at the association of medication with relapse adjusted for each covariate separately to investigate confounding of the association with medication and relapse. Following a post-hoc test for collinearity between smoking and harmful substance use we ran the multivariable model excluding harmful substance use. Since women on no medication may be non-adherent or psychiatrically stable, we ran a test for the interaction of psychiatric stability (admission in the 2 years before pregnancy) and medication on relapse. Most research has investigated bipolar disorder so we also tested the interaction of medication with diagnosis on relapse and ran the analysis stratified by diagnostic group (affective/ affective), reclassifying medication into two categories – continued versus stopped/switched/no medication. All interaction tests used the likelihood ratio.

A number of sensitivity analyses were carried out. To address issues of reverse causality for medication change and relapse both occurring in the first trimester, pregnancies with first trimester relapses were excluded, followed by an evaluation, censoring pregnancies which ended in the 2<sup>nd</sup> trimester. For comparability with other investigations of admissions in pregnancy, a sensitivity analysis excluded self-harm from the definition of relapse. To compare with previous studies on stopping medication and stopping abruptly in pregnancy (Viguera et al, 2007), women who switched a medication only were excluded from the 'stopped or switched' group (statistical power was insufficient to analyse switching separately) and we further excluded those stopping medication gradually. A further analysis excluded women with possible non-adherence in case of

misclassification and we also reclassified women who continued a medication but were coded non-adherent into the stopped/switched group. Finally we looked at the unadjusted and adjusted associations with stopping versus switching/ continuing an antidepressant, mood stabiliser and antipsychotic separately.

## **Results**

### **Sample characteristics**

There were 454 pregnancies in 389 women with history of SMI (Figure 1). Fifty-two (13.4%) women had more than one pregnancy in the study period (number of pregnancies per woman ranged from 1-5). Table 1 displays demographic and clinical characteristics of women. Of 203 (52.2%) women with diagnoses of non-affective SMI, 112 (28.8%) had diagnoses of schizophrenia or delusional disorder, 27 (6.9%) with schizoaffective disorder and 64 (16.5%) had other non-affective diagnoses including acute and transient psychosis; 186 (47.8%) women had affective diagnoses including bipolar disorder (138 women, 35.5%), depressive psychosis (41 women, 10.5%) and history of postpartum psychosis (7 women, 1.8%). Comorbid diagnoses between 9 months and 2 years before delivery episode included 18 (4.6%) women with substance use diagnoses, 9 (2.3%) with personality disorder diagnoses, 11 (2.8%) with anxiety disorders, 6 (1.5%) with learning difficulties and <5 with other diagnoses including eating disorders, conduct disorders and epilepsy.

In 454 pregnancies, there were 83 (18.3%) relapses, including 74 (16.3%) admissions to acute care (inpatient or home treatment), with 26 (5.7%) relapses occurring in the first trimester and 32 (7.5%) in the second. Thirteen pregnancies ended in the second trimester, and of 383 remaining pregnancies a further 25 (6.5%) relapsed in third trimester. All admissions to the mother and baby unit were for acute mental health reasons. Supplementary Table 1 provides further detail on relapses by trimester and diagnosis.

First trimester medication (antidepressant, mood stabiliser or antipsychotic) exposures were as follows: 147 (15.7%) no regular medication, 240 (52.9%) receiving an antipsychotic, 67 (14.8%) a mood stabiliser and 106 (23.4%) an antidepressant; 188 (41.4%) exposed to one agent only, 88

(19.3%) exposed to two and 31 (6.8%) to more than two. Table 1 shows medication categories in the 1<sup>st</sup> trimester as used for analysis. Medication categories were associated strongly from one trimester to the next; from 3 months before pregnancy to 1<sup>st</sup> trimester,  $\chi^2=312.64$ ,  $p<0.001$ , from 1<sup>st</sup> to 2<sup>nd</sup> trimester,  $\chi^2=296.41$ ,  $p<0.001$  and from 1<sup>st</sup> to 3<sup>rd</sup> trimester,  $\chi^2=182.34$ ,  $p<0.001$ .

Deprivation scores were missing for 18 (4.0%) pregnancies. HoNOS scores were missing for 212 (46.7%).

*Socio-demographic and clinical characteristics associated with relapse in pregnancy (Table 2) and associated with medication exposures (supplementary table 2).*

Relapse in pregnancy was significantly associated with non-white ethnicity, DV, smoking, non-affective psychosis, substance misuse, admissions in the two years before pregnancy, less than 2 years since previous acute admission, and self-harm in the 2 years before pregnancy. Very similar results were found when relapses in the 1st trimester were excluded. Women who continued to take medication had fewer relapses than women who were medication free or stopped/switched medication but this was not statistically significant. Age, history of admissions variables, self-harm and smoking in pregnancy were distributed differently across medication groups. Average age was highest in the continued medication group. Women with pregnancies unexposed to medication tended to have less admissions or time in acute care in the two years before pregnancy. Rates of self-harm were highest in those who stopped or switched a medication and lowest in those who continued. Recorded smoking was highest in those who stopped or switched medication and lowest in those unexposed to medication at the beginning of pregnancy.

Multivariable analysis

In the fully adjusted model (Table 3), (adjusted for medication group, admissions and self-harm in 2 years before pregnancy, diagnosis at baseline, harmful substance use, smoking in pregnancy and ethnicity) relapse was independently associated with non-affective psychosis, illness severity (admissions and self-harm in two years before pregnancy), substance misuse, smoking in pregnancy and non-white ethnicity. Given substance misuse and smoking were strongly associated ( $p < 0.001$ ), we also ran the model excluding substance misuse which made no difference to our findings. Women not on medication were more likely to relapse than women who continued medication; stopping or switching medication was not associated with increased risk of relapse (Table 3 & 4). However, when first trimester relapses were excluded, the association between being on no medication and relapse was attenuated (adjusted OR=1.43; 95%CI 0.69-2.98). Post-hoc analysis showed the association with relapse was most confounded by admissions in 2 years before pregnancy and smoking (supplementary table 3). Stratified by affective/ non-affective diagnosis (Supplementary Table 4), findings were similar, though due to smaller numbers most associations did not reach statistical significance. No significant interactions were found on risk of relapse between medication and affective/ non-affective diagnosis ( $\chi^2=0.01$ ,  $p=0.922$ ) and medication and admissions in the 2 years before pregnancy ( $\chi^2=0.52$ ,  $p=0.471$ ), or between DV and medication status ( $\chi^2=1.34$ ,  $p=0.248$ ).

### Sensitivity analyses

Defining relapse by acute admissions only ( $n=74$  relapses) made little difference to results. Excluding pregnancies where medication was stopped gradually ( $n=13$ ) made no meaningful difference (Table 4). Excluding those with recorded non-adherence to address potential misclassification of medication (17 pregnancies) strengthened the unadjusted difference between continuing vs. stopping/switching medication, but multivariable output remained similar (Supplementary Table 5). Exclusion of 14 pregnancies which ended in 2<sup>nd</sup> trimester made no difference (supplementary table

6). There was no association between these early ending pregnancies and relapse (Fisher's exact,  $p=0.482$ ). Women who stopped an antipsychotic had higher odds of relapse than women who continued/ switched, but this was not the case for mood stabilisers or antidepressants. However these findings were not significant (supplementary table 7).

## **Discussion**

In 389 women with SMI, there were 83 (18.3%) relapses during 454 pregnancies – 58 (24.4%) in pregnancies of women with non-affective diagnoses and 25 (11.6%) in women with affective disorders. Based on multivariable analyses, the nature and severity of the mental disorder (non-affective psychosis at baseline, recent admissions, harmful substance use, and recent self-harm) was associated with relapse in pregnancy. Smoking and non-white ethnicity also independently predicted relapse. Continuing medication compared with not taking medication at the beginning of pregnancy was protective, but we were unable to draw conclusions about switching or stopping medication in the first trimester.

These findings provide some reassurance that although relapse can occur in pregnancy, most women in this cohort did not have episodes needing acute care or manifesting as self-harm. Our findings suggest that risk of relapse in women with bipolar disorder may be lower than in studies of women attending specialist clinics (Viguera et al, 2011; Viguera et al, 2007), and comparable with the few studies carried out on more representative samples (Judd et al, 2014). There are few data in women with non-affective psychoses. The only other large study to date, which used administrative data, found that 24% of pregnant women with schizophrenia were admitted or visited a mental health emergency service, which is comparable to our findings (Vigod et al, 2014).

There has been little previous research into risk factors for relapse of severe mental disorder in pregnancy, although associations with severity of the pre-existing psychiatric illness have been reported (Munk-Olsen et al, 2009; Vigod & Ross, 2010; Viguera et al, 2007). Using comprehensive



clinical data we found that smoking, recent self-harm and substance misuse also showed associations with relapse, potentially suggesting higher-risk lifestyles. Such women may find it more difficult to prioritise stopping smoking or substance use in pregnancy; alternatively, smoking and substance misuse may be associated with worsening mental health. Both are important risk factors for adverse fetal outcomes (Stein et al, 2014), which need to be addressed by services. Being of black ethnicity was also associated with increased risk of relapse in pregnancy, consistent with research showing higher rates of relapse and worse outcomes in people of black compared with white ethnicity (Mann et al, 2014). Non-affective diagnosis has also been associated with worse outcomes in patients with psychosis (Morgan et al, 2014) as well as non-adherence to medication (Porcelli et al, 2016). Relapse or psychiatric admission in pregnancy may be an important predictor of relapse postpartum and further work with this cohort has looked at relapse in the first three months postpartum including admission in pregnancy as a predictor.

We did not find an association between medication discontinuation or switching and relapse, in contrast to others (Viguera et al, 2007) who found 86% risk in women with bipolar disorder stopping mood stabilising medication compared with 37% in women who continued, independent of disorder severity measures. Different definitions of relapse may explain some of this heterogeneity. In Viguera's study, it could not be determined whether medication change in the first trimester occurred before or after a relapse in early pregnancy; however, excluding women who relapsed in the first trimester attenuated our association, and indeed Viguera's data showed particularly high risk of relapse in early pregnancy. In observational research, independent effects of psychiatric illness and medication can be challenging to disentangle. Different classification of medication exposures may also explain some discrepancy in findings. Viguera investigated stopping mood stabilisers from 6 months before to 12 weeks after conception and excluded women who stopped medication over 6 months before. In our data women who stopped medication before pregnancy would be in the 'no medication' group and tended to relapse early.

To our knowledge, there are no comparative data in pregnant women with schizophrenia; however, outside the perinatal period, switching medication appears well tolerated (Roussidis et al, 2013).

These clinical records enabled us to investigate potential impact of other moderators, notably DV.

This needs replication in larger samples but its potential adverse impact on mental health is reported elsewhere (Khalifeh et al, 2015) as is the importance of addressing DV perinatally (NICE, 2014).

A major strength of this study was the novel methodology using electronic secondary mental health data. This allowed investigation of a larger cohort of women than previous clinical studies, with more detail on illness severity and other psychosocial characteristics than population registries.

SLaM provides near-monopoly secondary mental health services to its catchment, so these data are likely to include most pregnant women with SMI in this locality. The linkage with HES enabled us to collect psychiatric admissions data covering England, minimising loss to follow-up and giving a full picture of illness history for this dynamic cohort. We were also able to collect data on recorded DV, substance use and self-harm. Detailed clinical data on medication enabled us to capture changes resulting from mutual decisions made under clinical guidance (Swanson et al, 2015) or self-initiated cessation decisions, as well as recorded adherence, so we were able to address some misclassifications not achievable with registry data. CRIS data are not collected for research and rely on accuracy and comprehensiveness of clinical note keeping. Although we used multiple validated search terms, some information may be absent or underreported, such as that on smoking. We were not able to ascertain previous history of perinatal episodes which have been shown predictive of postpartum relapse. However, this methodology enabled us to observe this high-risk group of pregnant women in a real-world setting. We were able to include self-harm and women potentially more severely at risk than may take part in clinical studies.

Considering sample selection, our data source does not include women managed only in primary care. However, the threshold for referral to secondary care may be lower in pregnancy, and national guidelines (NICE, 2014) recommend referral of pregnant women with SMI history to secondary

mental health care. There may also be selection into pregnancy in women with SMI, since reported fertility is lower than that in the general population (Howard et al, 2002; Vigod et al, 2012).

Schizophrenia may affect relationship stability (Howard, 2005) and there are also reported higher rates of termination (Laursen & Munk-Olsen, 2010) and unplanned pregnancies in women with SMI (Miller & Finnerty, 1996). These factors affect comparability with times outside pregnancy. The linkage to HES data allowed a robust method of identifying pregnancies, covering deliveries across England. Home births are thought to be under recorded in HES (2.4% of deliveries in England in 2011 were at home) (ONS, 2013) though pregnancies in women with SMI are considered high-risk obstetrically so home births may be less likely. The retrospective design may have identified a particularly severe group of women since by default women admitted in pregnancy would have been included in the cohort. Women also receive many diagnoses in the health records; therefore, potential misclassification may have affected inclusion.

Considering the outcome, admission to acute care may not be a sensitive indicator of relapse, as less severe relapses may be managed by the secondary services already providing care, or by medication changes. Whilst using this endpoint may have missed some cases of relapse, it may also have included women admitted for social reasons, thus reflecting socio-economic disadvantage rather than relapse *per se*. However, all admissions to the MBU during pregnancy were for acute illness so this is unlikely to have skewed the results significantly. We did not carry out survival analyses as dates of admission were considered to be insufficiently accurate estimates of end-point timing.

Considering exposures, the stopped/switched medication group may be heterogeneous in terms of reasons for stopping/switching. However most women who stopped medication in the first trimester did so because of pregnancy (Taylor et al, 2015) and sensitivity analyses excluding switchers and gradual stoppers made little difference to findings. We did not have the power to analyse gradual stoppers separately and there may have been some misclassification of speed of stopping medication as it was difficult to determine from the clinical notes and we assumed abrupt

stopping unless stated otherwise. Previous research has shown that women with bipolar disorder may have higher risk of relapse if they discontinue or do not take medication. Women with non-affective psychosis may not be the same. However we did not have a large enough sample to ascertain predictors or moderators of relapse separately in the non-affective and affective groups. The fairly small proportion of relapses detected limited statistical power to detect some associations, and larger cohorts would be needed to address some remaining questions. Prospective clinical research could address some of the issues of unrecorded clinical information although assembling a large cohort would be costly and time-consuming with higher risk of attrition bias.

### Conclusions

Women with indicators of more severe SMI are more likely to relapse in pregnancy, these indicators including having more admissions in the two years prior to pregnancy, non-affective illness, history of self-harm, smoking and substance use. We found weaker evidence of association with medication change but women on no medication at the start of pregnancy were more likely to relapse in early pregnancy. While the risks and benefits of taking medication in pregnancy must be weighed up for each individual (NICE, 2014) this study has identified those at particularly high risk of relapse in pregnancy – evidence that could be shared with women who are making decisions about medication in pregnancy. Women with SMI have multiple risk factors for adverse pregnancy outcomes including risk of relapse, smoking, substance use, medication and should be under care of specialist services during pregnancy. Attention should also be paid to potentially modifiable risk factors including smoking and substance use.

**Financial support:** LMH and CT are supported by an NIHR Research Professorship (NIHR-RP-R3-12-011). RS & MB are part-funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. This study has been part-funded by the Johnson & Johnson CSR Citizenship Trust. Johnson & Johnson had no involvement in the design of the study, or access to any data throughout the

project. This research was also supported by the Biomedical Research Nucleus data management and informatics facility at South London and Maudsley NHS Foundation Trust, which is funded by the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and a joint infrastructure grant from Guy's and St Thomas' Charity and the Maudsley Charity.

## **References**

*Health & Social Care Information Centre* Available online:

<http://systems.hscic.gov.uk/data/clinicalcoding/codingstandards/opcs4> [Accessed.

Austin, M., Kildea, S. & Sullivan, E. (2007) Maternal mortality and psychiatric morbidity in the perinatal period: challenges and opportunities for prevention in the Australian setting. *Medical Journal of Australia*, 186(7), 364.

Bursac, Z., Gauss, C. H., Williams, D. K. & Hosmer, D. W. (2008) Purposeful selection of variables in logistic regression. *Source code for biology and medicine*, 3(1), 1.

Cantwell, R., Clutton-Brock, T., Cooper, G., Dawson, A., Drife, J., Garrod, D., Harper, A., Hulbert, D., Lucas, S. & McClure, J. (2011) Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006-2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG: an international journal of obstetrics and gynaecology*, 118, 1-205.

Cunningham, H., Tablan, V., Roberts, A. & Bontcheva, K. (2013) Getting more out of biomedical documents with GATE's full lifecycle open source text analytics. *PLoS computational biology*, 9(2), e1002854.

Di Florio, A., Forty, L., Gordon-Smith, K., Heron, J., Jones, L., Craddock, N. & Jones, I. (2013) Perinatal Episodes Across the Mood Disorder Spectrum. *Jama Psychiatry*, 70(2), 168-175.

Fernandes, A. C., Cloete, D., Broadbent, M. T., Hayes, R. D., Chang, C. K., Jackson, R. G., Roberts, A., Tsang, J., Soncul, M., Liebscher, J., Stewart, R. & Callard, F. (2013) Development and evaluation of a

de-identification procedure for a case register sourced from mental health electronic records. *BMC Med Inform Decis Mak*, 13, 71.

Grof, P., Robbins, W., Alda, M., Berghoefer, A., Vojtechovsky, M., Nilsson, A. & Robertson, C. (2000) Protective effect of pregnancy in women with lithium-responsive bipolar disorder. *Journal of Affective Disorders*, 61(1-2), 31-9.

Howard, L. M. (2005) Fertility and pregnancy in women with psychotic disorders. *Eur J Obstet Gynecol Reprod Biol*, 119(1), 3-10.

Howard, L. M., Kumar, C., Leese, M. & Thornicroft, G. (2002) The general fertility rate in women with psychotic disorders. *American Journal of Psychiatry*, 159(6), 991-7.

Johnson, S., Needle, J., Bindman, J. P. & Thornicroft, G. (2008) *Crisis resolution and home treatment in mental health*. New York: Cambridge University Press.

Jones, I., Chandra, P. S., Dazzan, P. & Howard, L. M. (2014) Bipolar disorder, affective psychosis, and schizophrenia in pregnancy and the post-partum period. *The Lancet*, 384(9956), 1789-1799.

Judd, F., Komiti, A., Sheehan, P., Newman, L., Castle, D. & Everall, I. (2014) Adverse obstetric and neonatal outcomes in women with severe mental illness: To what extent can they be prevented? *Schizophrenia research*, 157(1), 305-309.

Kadra, G., Stewart, R., Shetty, H., Jackson, R. G., Greenwood, M. A., Roberts, A., Chang, C.-K., MacCabe, J. H. & Hayes, R. D. (2015) Extracting antipsychotic polypharmacy data from electronic health records: developing and evaluating a novel process. *BMC psychiatry*, 15(1), 1.

Khalifeh, H., Hunt, I. M., Appleby, L. & Howard, L. M. (2016) Suicide in perinatal and non-perinatal women in contact with psychiatric services: 15 year findings from a UK national inquiry. *The Lancet Psychiatry*.

Khalifeh, H., Moran, P., Borschmann, R., Dean, K., Hart, C., Hogg, J., Osborn, D., Johnson, S. & Howard, L. (2015) Domestic and sexual violence against patients with severe mental illness. *Psychological medicine*, 45(04), 875-886.

- Laursen, T. M. & Munk-Olsen, T. (2010) Reproductive patterns in psychotic patients. *Schizophrenia research*, 121(1), 234-240.
- Li, Q., Andrade, S. E., Cooper, W. O., Davis, R. L., Dublin, S., Hammad, T. A., Pawloski, P. A., Pinheiro, S. P., Raebel, M. A., Scott, P. E., Smith, D. H., Dashevsky, I., Haffnerreffer, K., Johnson, K. E. & Toh, S. (2013) Validation of an algorithm to estimate gestational age in electronic health plan databases. *Pharmacoepidemiology and drug safety*, 22(5), 524-32.
- Mann, F., Fisher, H. L., Major, B., Lawrence, J., Tapfumaneyi, A., Joyce, J., Hinton, M. F. & Johnson, S. (2014) Ethnic variations in compulsory detention and hospital admission for psychosis across four UK Early Intervention Services. *BMC Psychiatry*, 14(1), 256.
- Miller, L. J. & Finnerty, M. (1996) Sexuality, pregnancy, and childrearing among women with schizophrenia-spectrum disorders. *Families and Mental Health Treatment: A Compendium of Articles from Psychiatric Services and Hospital and Community Psychiatry*, 45.
- Morgan, C., Lappin, J., Heslin, M., Donoghue, K., Lomas, B., Reininghaus, U., Onyejiaka, A., Croudace, T., Jones, P. B. & Murray, R. M. (2014) Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. *Psychological medicine*, 44(13), 2713-2726.
- Munk-Olsen, T., Laursen, T. M., Mendelson, T., Pedersen, C. B., Mors, O. & Mortensen, P. B. (2009) Risks and predictors of readmission for a mental disorder during the postpartum period. *Arch Gen Psychiatry*, 66(2), 189-95.
- NICE (2014) *Antenatal and postnatal mental health*, 2014. Available online: <http://www.nice.org.uk/guidance/cg192> [Accessed.
- Noble, M., McLennan, D., Wilkinson, K., Whitworth, A., Exley, S., Barnes, H. & Dibben, C. (2007) *The English indices of deprivation*. London.
- ONS (2013) *Births in England and Wales by Characteristics of Birth 2, 2011*, 2013. Available online: [http://www.ons.gov.uk/ons/dcp171778\\_298892.pdf](http://www.ons.gov.uk/ons/dcp171778_298892.pdf) [Accessed.
- Perera, G., Broadbent, M., Callard, F., Chang, C.-K., Downs, J., Dutta, R., Fernandes, A., Hayes, R. D., Henderson, M. & Jackson, R. (2016) Cohort profile of the South London and Maudsley NHS

Foundation Trust Biomedical Research Centre (SLaM BRC) Case Register: current status and recent enhancement of an Electronic Mental Health Record-derived data resource. *BMJ open*, 6(3), e008721.

Polling, C., Tulloch, A., Banerjee, S., Cross, S., Dutta, R., Wood, D., Dargan, P. & Hotopf, M. (2015) Using routine clinical and administrative data to produce a dataset of attendances at Emergency Departments following self-harm. *BMC emergency medicine*, 15(1), 15.

Porcelli, S., Bianchini, O., De Girolamo, G., Aguglia, E., Crea, L. & Serretti, A. (2016) Clinical factors related to schizophrenia relapse. *International journal of psychiatry in clinical practice*, 20(2), 54-69.

Rochon-Terry, G., Gruneir, A., Seeman, M. V., Ray, J. G., Rochon, P., Dennis, C.-L., Grigoriadis, S., Fung, K., Kurdyak, P. A. & Vigod, S. N. (2016) Hospitalizations and Emergency Department Visits for Psychiatric Illness During and After Pregnancy Among Women With Schizophrenia. *The Journal of clinical psychiatry*.

Roussidis, A., Kalkavoura, C., Dimelis, D., Theodorou, A., Ioannidou, I., Mellos, E., Mylonaki, T., Spyropoulou, A. & Yfantis, A. (2013) Reasons and clinical outcomes of antipsychotic treatment switch in outpatients with schizophrenia in real-life clinical settings: the ETOS observational study. *Annals of general psychiatry*, 12(1), 42.

Stein, A., Pearson, R. M., Goodman, S. H., Rapa, E., Rahman, A., McCallum, M., Howard, L. M. & Pariante, C. M. (2014) Effects of perinatal mental disorders on the fetus and child. *The Lancet*, 384(9956), 1800-1819.

Stewart, R., Soremekun, M., Perera, G., Broadbent, M., Callard, F., Denis, M., Hotopf, M., Thornicroft, G. & Lovestone, S. (2009) The South London and Maudsley NHS foundation trust biomedical research centre (SLAM BRC) case register: development and descriptive data. *BMC Psychiatry*, 9(1), 51.

Swanson, S. A., Hernandez-Diaz, S., Palmsten, K., Mogun, H., Olfson, M. & Huybrechts, K. F. (2015) Methodological considerations in assessing the effectiveness of antidepressant medication



continuation during pregnancy using administrative data. *Pharmacoepidemiology and drug safety*, 24(9), 934-942.

Taylor, C. L., Stewart, R., Ogden, J., Broadbent, M., Pasupathy, D. & Howard, L. M. (2015) The characteristics and health needs of pregnant women with schizophrenia compared with bipolar disorder and affective psychoses. *BMC Psychiatry*, 15, 88.

Taylor, C. L., van Ravesteyn, L. M., Lambregtse van denBerg, M. P., Stewart, R. J. & Howard, L. M. (2016) The prevalence and correlates of self-harm in pregnant women with psychotic disorder and bipolar disorder. *Archives of women's mental health*, 1-7.

Vigod, S., Kurdyak, P., Dennis, C., Gruneir, A., Newman, A., Seeman, M., Rochon, P., Anderson, G., Grigoriadis, S. & Ray, J. (2014) Maternal and newborn outcomes among women with schizophrenia: a retrospective population-based cohort study. *BJOG: An International Journal of Obstetrics & Gynaecology*, 121(5), 566-574.

Vigod, S. N. & Ross, L. E. (2010) Epidemiology of psychotic symptoms during pregnancy and postpartum in women with schizophrenia. *Current Women's Health Reviews*, 6(1), 17-21.

Vigod, S. N., Seeman, M. V., Ray, J. G., Anderson, G. M., Dennis, C. L., Grigoriadis, S., Gruneir, A., Kurdyak, P. A. & Rochon, P. A. (2012) Temporal trends in general and age-specific fertility rates among women with schizophrenia (1996-2009): a population-based study in Ontario, Canada. *Schizophr Res*, 139(1-3), 169-75.

Viguera, A. C., Tondo, L., Koukopoulos, A. E., Reginaldi, D., Lepri, B. & Baldessarini, R. J. (2011) Episodes of mood disorders in 2,252 pregnancies and postpartum periods. *Am J Psychiatry*, 168(11), 1179-85.

Viguera, A. C., Whitfield, T., Baldessarini, R. J., Newport, D. J., Stowe, Z., Reminick, A., Zurick, A. & Cohen, L. S. (2007) Risk of recurrence in women with bipolar disorder during pregnancy: prospective study of mood stabilizer discontinuation. *American Journal of Psychiatry*, 164(12), 1817-24; quiz 1923.

Wesseloo, R., Kamperman, A. M., Munk-Olsen, T., Pop, V. J., Kushner, S. A. & Bergink, V. (2015) Risk of postpartum relapse in bipolar disorder and postpartum psychosis: a systematic review and meta-analysis. *American Journal of Psychiatry*.

Wing, J., Beevor, A., Curtis, R., Park, S., Hadden, S. & Burns, A. (1998) Health of the Nation Outcome Scales (HoNOS). Research and development. *The British Journal of Psychiatry*, 172(1), 11-18.

Wu, C.-Y., Chang, C.-K., Robson, D., Jackson, R., Chen, S.-J., Hayes, R. D. & Stewart, R. (2013) Evaluation of smoking status identification using electronic health records and open-text information in a large mental health case register. *PloS one*, 8(9), e74262.

Table 1: Socio-demographic and clinical characteristics of 389 pregnant women with severe mental illness

	N=389
<b>Diagnosis at beginning of pregnancy, N(%)</b>	
Affective	186 (47.8)
Non affective	203 (52.2)
<b>Ethnicity, N(%)</b>	
White British & other White	132 (33.9)
Black African & other Black	188 (48.3)
Asian/ Mixed/ Other	69 (17.7)
<b>Deprivation score, median(range)<sup>1</sup></b>	34.9 (4.8,61.5)
<b>Maternal age at 1<sup>st</sup> index delivery, mean(SD),</b>	31.8 (6.1)
<b>Partner during 1<sup>st</sup> index pregnancy</b>	269 (69.2)
<b>Number of children at 1<sup>st</sup> index pregnancy, N(%)</b>	
0	182 (46.8)
1	103 (26.5)
≥2	104 (26.7)
<b>Recorded history of child abuse, N(%)</b>	78 (20.1)
<b>Recorded domestic abuse before/during pregnancy, N(%)</b>	133 (34.2)
<b>Smoking in pregnancy, N(%)</b>	69 (17.7)
<b>Harmful substance use, N(%)</b>	96 (24.7)
<b>Self-harm in 2 years before pregnancy, N(%)</b>	54 (13.9)
<b>Number of days of acute care in 2 years before pregnancy, N(%)</b>	
0	235 (60.4)
1-30	53 (13.6)
31-60	43 (11.1)
61+ (-537)	58 (14.9)
<b>Number of acute admissions in 2 years before pregnancy, N(%)</b>	
0	235 (60.4)
1	103 (26.5)
2	30 (7.7)
>2	21 (5.4)
<b>Time since last acute admission, N(%)</b>	
None in 2 years before pregnancy	235 (60.4)
1 year	79 (20.3)
2 years	75 (19.3)
<b>MHA<sup>3</sup> in 2 years before pregnancy, N(%)</b>	67 (17.2)
<b>Highest HoNOS total adjusted score in 2 years before pregnancy, median (range)<sup>2</sup></b>	12 (0-36)
<b>Family history of any mental illness, N(%)</b>	168 (43.2)
<b>Medication changes 1<sup>st</sup> trimester, N(%)</b>	

Continued	145 (37.3)
Stopped and/or switched	105 (27.0)
No medication at start	139 (35.7)

1: n=374, 2: n=196, 3: detained under section 2 or 3 of the mental health act

Table 2 – Socio-demographic and clinical characteristic, and associations with relapse in pregnancy episodes (n=454)

	N	N(%) relapse in pregnancy (n=83)	Odds ratio (95% CI)	p-value
<b>Baseline Diagnosis</b>				
Affective	216	25 (11.6)	<b>Ref</b>	
Non affective	238	58 (24.4)	2.58 (1.45,4.58)	0.001*
<b>Ethnicity</b>				
White British & other White	221	49 (22.2)	<b>Ref</b>	
Black African & other Black	154	17 (11.0)	2.50 (1.24,5.04)	0.010*
Asian/ Mixed/ Other	79	17 (21.5)	2.31 (1.01,5.25)	0.047*
<b>Partner during pregnancy<sup>2</sup></b>	321	58 (18.1)	0.94 (0.53,1.65)	0.826
<b>Children at index pregnancy</b>				
0	188	35 (18.6)	<b>Ref</b>	
1	127	25 (19.7)	1.06 (0.58,1.98)	0.836
≥2	139	23 (16.6)	0.83 (0.44,1.57)	0.563
<b>Victim of child abuse<sup>2</sup></b>	96	22 (22.9)	1.44 (0.80,2.60)	0.221
<b>Victim of domestic abuse before or during pregnancy<sup>2</sup></b>	167	39 (23.4)	1.74 (1.02,2.98)	0.044*
<b>Smoking in pregnancy<sup>2</sup></b>	79	31 (39.2)	4.47 (2.23,8.99)	<0.001*
<b>Harmful substance use<sup>2</sup></b>	109	36 (33.0)	3.37 (1.82,6.21)	<0.001*
<b>Self-harm in 2 years before pregnancy<sup>2</sup></b>	61	23 (37.7)	3.60 (1.83,7.10)*	<0.001*
<b>Number of days of acute care in 2 years before pregnancy,</b>				
0	270	38 (14.1)	<b>Ref</b>	
1-30	65	18 (27.7)	2.34 (1.22,4.48)*	0.010*
31-60	46	8 (17.4)	1.28 (0.56,2.97) <sup>a</sup>	0.558*

61+	73	19 (26.0)	2.15 (1.15,4.02)	0.017
<i>Linear association</i>	454	83 (18.3)	1.01 (1.00,1.02)*	0.091
<b>Number of acute admissions in 2 years before pregnancy</b>				
0	270	38 (14.1)	Ref	
1	121	25 (20.7)	1.59 (0.91,2.79)	0.107
2	39	11 (28.2)	2.40 (1.10,5.26)*	0.029*
>2	24	9 (37.5)	3.67 (1.47,9.25)*	0.005*
<i>Linear association</i>	454	83 (18.3)	1.55 (1.20,2.01)*	<0.001*
<b>MHA<sup>1</sup> 2 years before pregnancy<sup>2</sup></b>	77	16 (20.8)	1.18 (0.61,2.28)	0.663
<b>Time since last admission (years)</b>				
Not in 2 years before pregnancy	270	38 (14.1)	Ref	
<1 year	99	30 (30.3)	2.66 (1.53,4.63)	0.001*
1-2 years	85	15 (17.7)	1.31 (0.68,2.52)	0.425
<b>Family history of mental illness<sup>2</sup></b>	199	40 (20.1)	1.23 (0.74,2.04)	0.432
<b>Medication change 1<sup>st</sup> trimester</b>				
Continued/ no stop or switch	173	25 (14.5)	Ref	
Stopped and/ or Switched	118	25 (21.2)	1.66 (0.85,3.26)	0.140
No medication	163	33 (20.3)	1.60 (0.84,3.06)	0.153
<b>Deprivation score (per unit increase), (n=436)</b>			1.02 (0.99,1.04)	0.147
<b>Maternal age at delivery (per year increase), (n=454)</b>			0.96 (0.92,1.01)	0.091
<b>Highest HoNOS total adjusted score in 2 years before pregnancy (per unit increase), n=242</b>			1.01 (0.97,1.06)	0.599

<sup>1</sup>: detained under section 2 or 3 of the Mental Health Act; \*p<0.05

<sup>2</sup>: for partner during pregnancy, victim of child/ domestic abuse, smoking/ harmful substance use, MHA admission and family history of mental illness variables, the reference group was 'no' and 'absent information' combined

Table 3: Multivariable analysis of predictors of relapse in pregnancy

	Relapse in pregnancy, odds ratios (95% CI); N=454, n=83 relapses		Relapse in 2 <sup>nd</sup> or 3 <sup>rd</sup> trimester, odds ratios (95% CI); N=428, n=57 relapses	
	Unadjusted	Fully adjusted	Unadjusted	Fully adjusted
Medication				
Continuation	Ref	Ref	Ref	Ref
Stopped/ switched	1.66 (0.85,3.26), 0.140	1.30 (0.66,2.58), 0.446	1.41 (0.67,2.95), 0.363	1.16 (0.55,2.45), 0.703
No medication	1.60 (0.84,3.06), 0.153	<b>1.99 (1.05,3.75), 0.034</b>	1.01 (0.49,2.09), 0.977	1.43 (0.69,2.98), 0.335
Number of admissions in last 2 years	<b>1.55 (1.20,2.01), 0.001</b>	<b>1.37 (1.03,1.84), 0.033</b>	<b>1.68 (1.26,2.23), &lt;0.001</b>	<b>1.49 (1.08,2.05), 0.015</b>
Non-affective diagnosis at baseline	<b>2.58 (1.45,4.58), 0.001</b>	<b>2.03 (1.16,3.54), 0.013</b>	<b>2.55 (1.35,4.81), 0.004</b>	<b>2.06 (1.08,3.94), 0.028</b>
Self-harm in last 2 years	<b>3.60 (1.83,7.10), &lt;0.001</b>	<b>2.24 (1.15,4.34), 0.017</b>	<b>3.25 (1.38,7.66), 0.007</b>	1.87 (0.87,4.03), 0.109
Harmful Substance use	<b>3.37 (1.82,6.21), &lt;0.001</b>	<b>2.15 (1.13,4.08), 0.019</b>	<b>2.83 (1.37,5.83), 0.005</b>	1.59 (0.74,3.41), 0.231
Smoking in pregnancy	<b>4.47 (2.23,8.99), &lt;0.001</b>	<b>2.52 (1.26,5.02), 0.009</b>	<b>4.11 (1.83,9.22), 0.001</b>	<b>2.72 (1.21,6.09), 0.015</b>
Ethnicity				
White	Ref	Ref	Ref	Ref
Black	<b>2.50 (1.24,5.04), 0.010</b>	<b>2.37 (1.23,4.57), 0.010</b>	<b>2.92 (1.31,6.50), 0.009</b>	<b>2.79 (1.26,6.18), 0.014</b>
Mixed/ Other	<b>2.31 (1.01,5.25), 0.047</b>	<b>2.94 (1.32,6.56), 0.008</b>	<b>2.65 (1.04,6.77), 0.042</b>	<b>3.29 (1.27,8.53), 0.014</b>

Based on predictors with  $p < 0.2$  from bivariate analysis. Maternal age at delivery, deprivation score and domestic abuse were removed from the final model as they had a fully adjusted  $p > 0.5$

Table 4: Sensitivity analyses to investigate stopping medication in 1<sup>st</sup> trimester

	Excluding pregnancies with only a switch in medication, odds ratios (95% CI); N=436, n=79 relapses		Stopping medication abruptly (excluding only switchers and gradual stoppers), odds ratios (95% CI); N=422, n=75 relapses	
	Unadjusted	Fully adjusted	Unadjusted	Fully adjusted
Medication				
<i>Continued</i>	Ref	Ref	Ref	Ref
<i>Stopped</i>	1.65 (0.81,3.37), 0.170	1.30 (0.63,2.68), 0.475	1.48 (0.70,3.11), 0.302	1.16 (0.53,2.53), 0.704
<i>No medication at start</i>	1.61 (0.84,3.10), 0.152	<b>1.99 (1.03,3.84), 0.040</b>	1.62 (0.85,3.11), 0.146	<b>2.09 (1.06,4.13), 0.033</b>
Number of admissions in 2 years before	<b>1.52 (1.16,2.00), 0.003</b>	<b>1.39 (1.03,1.88), 0.034</b>	<b>1.52 (1.16,2.00), 0.003</b>	<b>1.40 (1.02,1.93), 0.038</b>
Baseline diagnosis	<b>2.56 (1.41,4.63), 0.002</b>	<b>1.99 (1.12,3.53), 0.019</b>	<b>2.56 (1.41,4.63), 0.002</b>	<b>2.01 (1.10,3.68), 0.023</b>
Self-harm in 2 years before pregnancy	<b>3.01 (1.49,6.08), 0.002</b>	<b>2.12 (1.05,4.28), 0.039</b>	<b>3.01 (1.49,6.08), 0.002</b>	<b>2.08 (1.00,4.35), 0.051</b>
Harmful substance use	<b>2.95 (1.57,5.54), 0.001</b>	<b>2.01 (1.02,3.95), 0.042</b>	<b>2.95 (1.57,5.54), 0.001</b>	1.97 (0.96,4.03), 0.063
Smoking	<b>4.28 (2.06,8.89), &lt;0.001</b>	<b>2.44 (1.16,5.10), 0.018</b>	<b>4.28 (2.06,8.89), &lt;0.001</b>	<b>2.75 (1.24,6.09), 0.012</b>
Ethnicity				
<i>White</i>	Ref	Ref	Ref	Ref
<i>Black</i>	<b>3.16 (1.43,6.98), 0.022</b>	<b>2.54 (1.20,5.38), 0.015</b>	<b>4.18 (1.72,10.19), 0.002</b>	<b>3.40 (1.45,8.00), 0.005</b>
<i>Mixed/ Other</i>	<b>2.84 (1.16,6.96), 0.005</b>	<b>3.12 (1.33,7.35), 0.009</b>	<b>3.52 (1.33,9.34), 0.012</b>	<b>4.21 (1.61,11.00), 0.003</b>

Based on predictors with  $p < 0.2$  from bivariate analysis. Maternal age at delivery, deprivation score and domestic abuse were removed from the final model as they had a fully adjusted  $p > 0.5$



